

Hot accretion discs with thermal Comptonisation and advection in luminous black hole sources

Andrzej A. Zdziarski

N. Copernicus Astronomical Center, Bartycka 18, 00-716 Warsaw, Poland

Accepted 1998 March 18. Received 1997 October 7

ABSTRACT

We solve for the structure of a hot accretion disc with unsaturated thermal Comptonisation of soft photons and with advection, generalizing the classical model of Shapiro et al. The upper limit on the accretion rate due to advection constrains the luminosity to $\lesssim 0.15y^{3/5}\alpha^{7/5}$ of the Eddington limit, where y and α are the Compton and viscosity parameters, respectively. The characteristic temperature and Thomson optical depth of the inner flow at accretion rates within an order of magnitude of that upper limit are $\sim 10^9$ K and ~ 1 , respectively. The resulting spectra are then in close agreement with the X-ray and soft γ -ray spectra from black-hole binaries in the hard state and Seyferts. At low accretion rates, bremsstrahlung becomes the dominant radiative process.

Key words: accretion: accretion discs – black hole physics – galaxies: Seyfert – gamma-rays: theory – X-rays: galaxies – X-rays: stars

1 INTRODUCTION

Matter accreting onto a compact object (either in an AGN or a binary) is likely to have a net angular momentum, which causes formation of a disc. Shakura & Sunyaev (1973) have developed a theory of optically-thick disc accretion. Such discs emit local radiation close to a blackbody, and thus their maximal temperature is at most $\sim 10^7$ K. Those discs cannot thus explain hard X-rays and soft γ -rays observed up to ~ 1 MeV from both black-hole binaries and Seyfert AGNs. Shapiro, Lightman & Eardley (1976, hereafter SLE76) found the existence of another branch of disc solution, in which the electron temperature, T_e , is $\sim 10^9$ K. In their specific model, the dominant radiative process is multiple Compton upscattering of copious soft photons by the disc electron plasma, which forms power-law X-ray spectra extending up to a few hundred of keV.

SLE76 assumed in their analysis that all the dissipated energy is locally radiated away. However, advection of hot ions is important in a range of parameters of the accretion flow (Ichimaru 1977). Solutions including both radiative cooling and advection have been extensively studied recently (e.g., Abramowicz et al. 1995; Narayan & Yi 1995, hereafter NY95; Chen et al. 1995; Björnsson et al. 1996). Some of those studies (e.g., Björnsson et al. 1996) assume Comptonised bremsstrahlung as the only cooling process, which turns out to reproduce well the qualitative properties of the solutions. However, the predicted X-ray spectra are then very hard, with the photon spectral index of $\Gamma \lesssim 1$, which disagrees with X-ray observations of both black-hole binaries

and Seyferts, showing typical $\Gamma \sim 1.7$ –2 (see, e.g., Zdziarski et al. 1997, hereafter Z97, for a review). Other studies (e.g., NY95, Narayan 1996) include also Comptonisation of thermal synchrotron photons in an equipartition magnetic field. However, as shown by SLE76, the synchrotron process does not supply sufficient flux of soft seed photons to explain the X-ray spectrum of Cyg X-1 in the hard state. This conclusion has been extended to GX 339-4 in the hard state and Seyferts by Zdziarski et al. (1998, hereafter Z98). Those results imply that if Comptonised synchrotron emission were the only source of soft photons, the X-ray spectra would be harder than those observed. On the other hand, soft-photon emission by a cold disc surrounding the hot flow has been shown by Esin (1997) either to be negligible or to lead to a runaway cooling of the hot flow.

In spite of these theoretical difficulties, X-ray observations show that a supply of soft seed photons sufficient to give rise to spectra with $\Gamma \sim 1.7$ –2 does exist in luminous black hole systems. Also, the shape of a thermal Comptonisation spectrum (at energies much above the seed photon energy) is a function solely of T_e and the Thomson optical depth, τ . Thus, it is possible to compare a model with X-ray data *without* specifying the soft photon source. Furthermore, most of the power in a spectrum with $\Gamma < 2$ is emitted at its high-energy end, and then even the model normalization can be compared with data without knowledge of the properties of soft seed photons.

Such an approach was adopted by SLE76, who specified the local cooling rate in a disc by assuming a constant value of the Compton parameter, y , a quantity directly related

to the X-ray index Γ . Observationally, luminous black-hole sources typically show an almost constant Γ with varying flux (e.g., Cyg X-1, Gierliński et al. 1997; GX 339-4, Ueda, Ebisawa & Done 1994; IC 4329A, Fiore et al. 1992), which justifies that assumption. The origin of soft seed photons does not need to be specified in the formalism, but SLE76 advocated its origin in cold clouds mixed with the hot flow (e.g., Kuncic, Celotti & Rees 1997). Thermal synchrotron radiation will also provide some soft photons.

In this work, we generalize the hot disc model of SLE76 by including advection, by treating the Compton parameter as a free parameter (whereas SLE76 assumed $y = 1$), and by extending the solution to low accretion rates. As we show below, the model predicts $\tau \sim 1$ and $T_e \sim 10^9$ K only weakly dependent on the disc radius and the accretion rate.

Thus, the predictions of the model are in excellent agreement with recent studies of broad-band ($\sim 1\text{--}10^3$ keV) spectra of both black-hole binaries in their hard state and Seyferts (e.g., Zdziarski, Johnson & Magdziarz 1996, Z97, Z98; Gierliński et al. 1997; Poutanen et al. 1997; Johnson et al. 1997; Magdziarz et al. 1998). Those studies show that the intrinsic spectra are well fitted by thermal Comptonisation in a plasma with just $T_e \sim 10^9$ K and $\tau \sim 1$, with a small dispersion of those parameters between different objects. Note that those parameters correspond to $y \lesssim 1$.

2 THE HOT DISC SOLUTION

In the model of SLE76, gravitational energy is converted into ion thermal energy, which is then transferred to electrons by Coulomb interactions. The electrons, in turn, radiate their energy by Compton upscattering the soft seed photons.

We assume here a pressure equipartition between the magnetic field and gas, $P_{\text{mag}} = P_{\text{gas}}$, which results in $P = 2(n_i kT_i + n_e kT_e)$, where T_i , n_i , T_e , n_e are the ion and electron temperature and number density, respectively. We then assume the plasma is two-temperature, i.e., $T_i \gg T_e$ (which is typically satisfied in an inner part of the disc), which allows us to neglect the electron contribution to P . The magnetic field here is treated similarly to, e.g., NY95, where the viscosity parameter, α , is unrelated to the magnetic field. In another model, Shakura & Sunyaev (1973) and Sakimoto & Coroniti (1981) use the magnetic field as a source of viscosity, which gives $P_{\text{mag}} \approx \alpha P_{\text{gas}}$. Adopting that model would slightly modify the equations below, slightly reducing τ for a given accretion rate. We further assume $e^\pm$ pairs are negligible (as shown to be the case in hot discs with advection by Björnsson et al. 1996 and Kusunose & Mineshige 1996), and neglect the radiation pressure.

We are mostly interested in an inner region of the disc, in which a self-similar solution of advection-dominated discs found by NY95 breaks down (e.g., Chen, Abramowicz & Lasota 1997). Therefore, we consider here local disc solutions and parametrize advection using results of global studies, following, e.g., Abramowicz et al. (1995) and Björnsson et al. (1996). As those authors, we use the pseudo-Newtonian potential (Paczynski & Wiita 1980), $\Phi = -GM/(R - R_S)$, and assume the disc rotation is Keplerian, $\Omega = (GM/R)^{1/2}/(R - R_S)$, where R is the disc radius, $R_S = 2GM/c^2$ is the Schwarzschild radius, and M is the central mass. We take into account the presence of He in the flow.

With the above assumptions, hydrostatic equilibrium and conservation of angular momentum yield (with the equation of state of $P = 2n_i kT_i$),

$$\left(\frac{H}{R}\right)^2 = \frac{2r\Theta_i q^2}{\mu_i}, \quad \Theta_i = \frac{J\mu_i \dot{m}}{2\alpha r^{3/2} g q \tau}, \quad (1)$$

respectively. Here H is the disc scale height, $r \equiv Rc^2/GM$, $q = 1 - 2/r$, $g = 1 + 4/3rq$, $J = 1 - (6/r)^{1/2}(3q/2)$ accounts for the zero-torque boundary condition at $r = 6$, $\Theta_i \equiv kT_i/m_p c^2$, α is the viscosity parameter, $\tau = n_e \sigma_T H$, $\dot{m} \equiv \dot{M} c^2/L_E$, $\dot{M}$ is the mass accretion rate, the Eddington luminosity is $L_E \equiv 4\pi\mu_e GM m_p c/\sigma_T$, $\mu_i = 4/(1 + 3X)$ and $\mu_e = 2/(1 + X)$ are the mean ion and electron molecular weight, respectively, $X (\approx 0.7)$ is the H mass fraction, and σ_T is the Thomson cross section.

The energy is dissipated in the disc at a rate of $Q = (3/4\pi)GM\dot{M}Jg/q^2R^3$ per unit area (including both sides of the disc). This energy is partly radiated and partly advected to smaller radii. We find that a quantity useful to describe the local emission of the disc is a local Eddington ratio, defined as the power per unit $\ln r$ in units of L_E , $\ell \equiv (dL/d \ln R)/L_E = 2\pi QR^2/L_E$. The dissipative ℓ is,

$$\ell = \frac{3Jg\dot{m}}{2q^2r} \equiv A\dot{m}, \quad (2)$$

where the function A is needed below. The maximum of $\ell(r)$ occurs at $r \approx 16.9$ (compared to $r = 27/2$ in the Newtonian, $q = g = 1$, $J = 1 - 6^{1/2}/r^{1/2}$, case). A fraction of ℓ is Coulomb-transferred to the electrons and then radiated. The rate of Coulomb transfer corresponds to (Spitzer 1956),

$$\ell_{\text{Coul}} = \frac{3}{(2\pi)^{1/2}} \frac{m_e f_C}{m_p \mu_i} \left(\frac{H}{R}\right)^{-1} \frac{r\tau^2}{\Theta_e^{3/2}} \left(\Theta_i - \frac{m_e}{m_p} \Theta_e\right). \quad (3)$$

Here $\Theta_e \equiv kT_e/m_e c^2$, $f_C \approx \ln \Lambda$, $\Lambda \approx 0.3\Theta_e(H/\tau a_0)^{1/2}$, a_0 is the Bohr radius, and $\Theta_i \ll \Theta_e \ll 1$ (which condition is typically satisfied in a hot flow). At this limit, the presence of He reduces the Coulomb rate with respect to the case of pure H by μ_e/μ_i . This is because the rate is proportional to Z^2/A_Z , which is unity for both H and He, and $n_i/n_e = \mu_e/\mu_i$. Here A_Z is the mass number of a Z th element. When the above condition is not satisfied, the rate given for H by Stepney & Guilbert (1983) can be generalized to yield an f_C of,

$$f_C = \ln \Lambda \frac{\pi^{1/2} \Theta_e^{3/2}}{2^{1/2} K_2(\Theta_e^{-1}) n_i} \sum_{Z=1} \frac{n_Z Z^2}{A_Z K_2(\Theta_Z^{-1})} \times \left[\frac{2(\Theta_e + \Theta_Z)^2 + 1}{\Theta_e + \Theta_Z} K_1\left(\frac{1}{\Theta_e} + \frac{1}{\Theta_Z}\right) + 2K_0\left(\frac{1}{\Theta_e} + \frac{1}{\Theta_Z}\right) \right], \quad (4)$$

where K_l is a modified Bessel function, n_Z is the density of the Z th element, and $\Theta_Z = \Theta_i/A_Z$. Note that NY95 neglected the effect of A_Z in their accounting for the presence of He in the Coulomb rate. Therefore, eq. (3.3) in NY95 and eq. (A1) in Mahadevan (1997) needs to be multiplied by 0.8 and $0.8\mu_e/\mu_i$, respectively, in the limit $\Theta_i \ll \Theta_e \ll 1$; otherwise eqs. (3)-(4) above should be used.

SLE76 assumed in their solution that a supply of soft seed photons results in the Compton parameter (e.g., Rybicki & Lightman 1979) of $y \equiv 4\Theta_e \tau \max(1, \tau) = 1$. This can be achieved, e.g., if most of the soft photons are from reprocessing of the hard radiation by cold matter in the vicinity of the hot flow (Haardt & Maraschi 1993). A given value of y

(which we allow here to differ from 1) approximately corresponds to a specific value of the energy amplification factor of Comptonisation, η (at constant soft photon energy, e.g., Dermer, Liang & Canfield 1991). Then, the local radiative balance can be expressed as,

$$\ell_{\text{Coul}} = \eta(y)\ell_{\text{soft}} + \ell_{\text{brem}}, \quad (5)$$

where ℓ_{soft} and ℓ_{brem} corresponds to the power in the soft photons and Comptonised bremsstrahlung, respectively. For now, we interpret ℓ_{Coul} as corresponding to Comptonisation of soft photons, which is equivalent to assuming $\ell_{\text{Coul}} \gg \ell_{\text{brem}}$. We also assume $T_i \gg T_e$, which allows us to neglect the negative term in eq. (3). At the end of this Section, we discuss the validity of those assumptions and discuss the behaviour of the solution when they break down.

We assume $\tau \lesssim 1$ (which we check it is the case *a posteriori*), which corresponds to T_e and ℓ_{Coul} of

$$\Theta_e = \frac{y}{4\tau}, \quad (6)$$

$$\ell_{\text{Coul}} = \frac{2^{3/2}3}{\pi^{1/2}} \frac{m_e}{m_p} \frac{f_C J^{1/2} \dot{m}^{1/2} \tau^3}{y^{3/2} \alpha^{1/2} q^{3/2} g^{1/2} r^{1/4}} \equiv B \dot{m}^{1/2} \tau^3. \quad (7)$$

A part of ℓ is advected to smaller radii (rather than radiated) at a rate of $Q_{\text{adv}} = (\dot{M}/2\pi R^2)(P/\mu_i n_i m_p)\xi$ (Chen & Taam 1993), where ξ is a dimensionless factor found ≈ 1 by global hot-disc studies (e.g., Chen et al. 1997). The advected power corresponds to,

$$\ell_{\text{adv}} = 2\dot{m}\Theta_e \xi \mu_i^{-1} = \frac{\xi J \dot{m}^2}{\alpha q g r^{3/2} \tau} \equiv C \dot{m}^2 \tau^{-1}. \quad (8)$$

The energy balance of the disc is given by,

$$\ell = \ell_{\text{Coul}} + \ell_{\text{adv}}. \quad (9)$$

Hereafter, we assume α , ξ , f_C and y to be constant; otherwise the solutions below of eq. (9) become implicit. The assumption of $y(r) = \text{constant}$ corresponds to a fixed ratio of the soft photon flux to the total local flux (eq. [5]), which can plausibly result from physical feedback, related to, e.g., geometry. The solution has two branches, corresponding to dominant cooling (SLE76), $\ell \approx \ell_{\text{Coul}} \propto \dot{m}$, and dominant advection, $\ell \approx \ell_{\text{adv}}$. The first one is,

$$\dot{m} = \frac{2^5 m_e^2 f_C^2 q r^{3/2} \tau^6}{\pi m_p^2 y^3 \alpha J g^3}, \quad \ell_{\text{Coul}}(r, \tau) = \frac{2^4 3 m_e^2 f_C^2 r^{1/2} \tau^6}{\pi m_p^2 y^3 \alpha q g^2}. \quad (10)$$

On the other hand, dominant advection corresponds to,

$$\dot{m} = \frac{3\alpha g^2 r^{1/2} \tau}{2\xi q}, \quad \ell_{\text{Coul}}(r, \tau) = \frac{3^{3/2} 2 m_e f_C (Jg)^{1/2} \tau^{7/2}}{\pi^{1/2} m_p y^{3/2} \xi^{1/2} q^2}, \quad (11)$$

$$\ell_{\text{Coul}}(r, \dot{m}) = \frac{2^{9/2} m_e}{3^2 \pi^{1/2} m_p} \frac{f_C \xi^3 J^{1/2} q^{3/2} \dot{m}^{7/2}}{\alpha^{7/2} y^{3/2} r^{7/4} g^{13/2}}. \quad (12)$$

Note no dependence on α in $\ell_{\text{Coul}}(r, \tau)$.

The general solution of eq. (9) is,

$$\tau = \frac{(A - Ca)^{2/5} a^{1/5}}{B^{2/5}} \quad \text{or} \quad \dot{m} = \frac{(A - Ca)^{2/5} a^{6/5}}{B^{2/5}}, \quad (13)$$

where the parameter $a \equiv \dot{m}/\tau$ is in the range $0 < a < 3\alpha g^2 r^{1/2}/2\xi q$ (with the lower and upper limits corresponding to dominant cooling and advection, respectively), and the functions A , B , C are defined in equations above. The local disc emission corresponds to,

$$\ell_{\text{Coul}} = (A - Ca)\dot{m} = \frac{(A - Ca)^{7/5} a^{6/5}}{B^{2/5}}. \quad (14)$$




Figure 1. Disc solutions at $r = 27/2$. The left-hand-side curves show the optically-thin solution for $\xi = 1$, $f_C = 15$. The disc emission is due to Comptonisation with $y = 1$ except for the solid curves below the break (at $\dot{m} \sim 0.002$ – 0.004), for which bremsstrahlung dominates. The right-hand-side curves show the optically-thick solution for $M = 10M_\odot$ and with a corona above the disc dissipating 1/2 of the total power (Svensson & Zdziarski 1994). Solid curves give the accreted $\dot{m}$ whereas the dashed curves give $\dot{m}_{\text{rad}}$, i.e., the part of $\dot{m}$ which is converted into escaping radiation. Note a good agreement of the $\dot{m}$ - τ dependence presented here with those of Abramowicz et al. (1995), Chen et al. (1995) and Björnsson et al. (1996).

Examples of the solution are shown in Fig. 1 (left-hand-side curves). Solid curves give $\dot{m}(\tau)$, and dashed curves give $\dot{m}_{\text{rad}}(\tau) \equiv \ell_{\text{Coul}}/A$, i.e., the locally-radiated fraction of $\dot{m}$ ($\dot{m}_{\text{rad}} = \dot{m}$ on the cooling branch). The corresponding temperature is given by eq. (6), $T_e \approx 1.5(y/\tau) \times 10^9$ K. The maxima of $\dot{m}$, $\dot{m}_{\text{rad}}$ and τ are located in the turning region connecting the cooling and advection branches at $a = 3A/4C$, $6A/13C$ and $A/3C$, respectively. The maximum $\dot{m}$ is,

$$\dot{m}_{\text{max}}(r) = \frac{3^{12/5} \pi^{1/5}}{2^{27/5}} \left(\frac{m_p}{m_e} \right)^{2/5} \frac{y^{3/5} \alpha^{7/5} J^{1/5} g^3 r^{3/10}}{f_C^{2/5} \xi^{6/5} q^{7/5}}. \quad (15)$$

Numerically, $\dot{m}_{\text{max}} \approx 2.85 y^{3/5} \alpha^{7/5} J^{1/5} g^3 r^{3/10}$, and, e.g., $\dot{m}_{\text{max}} \approx 7.37 y^{3/5} \alpha^{7/5}$ at $r = 27/2$. Hereafter, we set $\xi = 1$ and $f_C = 15$ (typical for stellar-mass black hole sources; for AGNs, $f_C \approx \ln \Lambda \approx 20$) in numerical expressions. The τ corresponding to $\dot{m}_{\text{max}}$ is given by

$$\tau(\dot{m}_{\max}) = \frac{3^{2/5} \pi^{1/5}}{2^{12/5}} \left(\frac{m_p}{m_e} \right)^{2/5} \frac{y^{3/5} \alpha^{2/5}}{\xi^{1/5} f_C^{2/5}} \frac{J^{1/5} g}{q^{2/5} r^{1/5}}. \quad (16)$$

Its maximum of $\tau(\dot{m}_{\max}) \approx 1.22 y^{3/5} \alpha^{2/5}$ is at $r = 12.9$. Advection at $\dot{m}_{\max}$ carries 0.59 of the dissipated energy, and the remainder corresponds to $\ell_{\text{Coul}}(\dot{m}_{\max})$ of,

$$\ell_{\text{Coul}}(\dot{m}_{\max}) = \frac{3^{17/5} \pi^{1/5}}{2^{42/5}} \left(\frac{m_p}{m_e} \right)^{2/5} \frac{y^{3/5} \alpha^{7/5} J^{6/5} g^4}{f_C^{2/5} \xi^{6/5} q^{17/5} r^{7/10}}, \quad (17)$$

or $\ell_{\text{Coul}}(\dot{m}_{\max}) \approx 1.07 y^{3/5} \alpha^{7/5} J^{6/5} g^4 q^{-17/5} r^{-7/10}$ numerically. Its maximum of $\ell_{\text{Coul}}(\dot{m}_{\max}) \approx 0.050 y^{3/5} \alpha^{7/5}$ is at $r = 18.6$. The maxima of τ and ℓ_{Coul} in the $\dot{m}$ - τ space (Fig. 1) are located in the cooling-dominated region and are larger than the values corresponding to $\dot{m}_{\max}$ by the factors of $(4/3)^{4/5} \approx 1.26$ and $2^{32/5} 7^{7/5} / 13^{13/5} \approx 1.63$, respectively.

We have assumed so far $T_i \gg T_e$, $\ell_{\text{Coul}} \gg \ell_{\text{brem}}$, where

$$\ell_{\text{brem}} = \frac{2^{5/2}}{\pi^{3/2}} \frac{m_e}{m_p} \alpha_f \left(\frac{H}{R} \right)^{-1} r \tau^2 A_{\text{brem}} \Theta_e^{1/2}, \quad (18)$$

$A_{\text{brem}} \geq 1$ is a factor including both Comptonisation and relativistic corrections and α_f is the fine-structure constant (Stepney & Guilbert 1983). (Note that the bremsstrahlung rate given in eq. [3.5] in NY95 appears incorrect: 1.25 should be replaced by μ_e .) The temperature ratio (at $\tau \lesssim 1$) and the relative bremsstrahlung rate (at $T_e \ll T_i$) are given by,

$$\frac{T_e}{T_i} = \frac{m_e \alpha y}{2 m_p \mu_i} \frac{g q r^{3/2}}{J \dot{m}}, \quad \frac{\ell_{\text{brem}}}{\ell_{\text{Coul}}} = \frac{\alpha_f A_{\text{brem}} \alpha y^2}{3 \pi f_C \tau} \frac{g q r^{3/2}}{J \dot{m}}. \quad (19)$$

We see that both ratios have very similar dependences on the disc parameters. On the advection branch, $\tau \propto \dot{m}$, and then $\ell_{\text{brem}}/\ell_{\text{Coul}} \propto \dot{m}^{-2}$, which implies bremsstrahlung becoming dominant at low $\dot{m}$ while still $T_e \ll T_i$. However, this happens only at very low ℓ_{Coul} (below the range of $\dot{m}_{\text{rad}}$ shown in Fig. 1). On the cooling branch, both above assumptions break down roughly at the same time. Note that the effects on $\tau(\dot{m})$ of the corrections due to $T_e \sim T_i$ and due to bremsstrahlung are in the opposite directions, thus partly canceling each other. Numerically, the $T_e \ll T_i$ condition (which is marginally more restrictive than $\ell_{\text{brem}} \ll \ell_{\text{Coul}}$) requires $\dot{m} \gg y \alpha / 20$ at $r = 27/2$, and $r \ll 280 / (\alpha y)^{2/3}$ at $\dot{m} = 1$. Otherwise, the disc is dominated by 1- T bremsstrahlung, for which the cooling-dominated solution is [for $P = 2(n_i + n_e)kT$, which modifies eq. (1)],

$$\dot{m} = \frac{2^3 \alpha_f}{3 \pi^{3/2}} \left[\frac{\mu_i \mu_e m_e}{(\mu_i + \mu_e) m_p} \right]^{1/2} A_{\text{brem}} \frac{q r^{3/2}}{g J} \tau^2. \quad (20)$$

From eqs. (1) and (6), we also find that a 1- T disc is characterized by a specific local y on both branches when $\tau < 1$,

$$y_{1T} = \frac{2 m_p \mu_e \mu_i J \dot{m}}{m_e (\mu_e + \mu_i) \alpha g q r^{3/2}}. \quad (21)$$

We see from the above discussion that our main assumptions: $T_e \ll T_i$, $\ell_{\text{brem}} \ll \ell_{\text{Coul}}$, and $y(r) = \text{constant}$, break down on the cooling branch roughly simultaneously at low $\dot{m}$ and/or large r . Thus, in Fig. 1, we show the 2- T , constant- y , solution at large $\dot{m}$ and the 1- T bremsstrahlung solution at small $\dot{m}$. In the latter, we set $A_{\text{brem}} = 1$ since both $y_{1T} \ll 1$ (eq. [21]) and $\Theta_e \ll 1$ (eq. [1]) at $\dot{m} \ll 1$. We simply assume a sharp transition between the two regimes; in reality, it will be smooth and more complex. We note that at radii at which bremsstrahlung dominates, $\dot{m}_{\max} \propto r^{-1/2}$ (Abramowicz et

al. 1995), whereas $\dot{m}_{\max} \propto r^{3/10}$ in the constant- y , 2- T , region (eq. [15]).

In contrast to the above results, Liang & Thompson (1979) and Wandel & Liang (1991) extended the 2- T solution of SLE76 (eq. [10]) to low values of $\dot{m}$ obtaining a cooling-dominated, 1- T solution with $y = \text{constant}$, $\dot{m} \propto \tau^{-1}$, $\tau > 1$. That extension neglects bremsstrahlung; however, we find the bremsstrahlung power on that solution exceeds the dissipated power, with the ratio of the two $\propto \tau^3$. Thus, that solution branch appears not to exist. Instead, the behaviour of the SLE76 solution at low $\dot{m}$ is given by a transition to the 1- T bremsstrahlung model, as shown above.

Note that y is necessarily variable in 1- T flows with $\tau < 1$ (eq. [21]) as well as it becomes large at small r , e.g., $y_{1T} \approx 7 \dot{m} / \alpha$ at $r = 27/2$. This represents a strong argument against their presence in inner regions of luminous black-hole systems, with constant $y < 1$ observed (Section 1).

3 DISCUSSION

We have solved the local structure of a hot accretion disc with the main cooling process being thermal Comptonisation of soft photons. Some feedback processes are assumed to lead to $y = \text{constant}$ in a range of radii, which corresponds to an approximately constant X-ray spectral index, Γ , of emission from each radius. The total spectrum can be then obtained by radial integration. The accretion rate of the hot flow can be constant throughout the flow, or, plausibly, it can equal the local maximum rate (eq. [15]). The latter case corresponds to a two-phase flow, in which a cold flow (e.g., in the form of clouds) carries the difference between the total $\dot{M}$ and that corresponding to the maximum $\dot{M}$ in the hot flow. The latter case maximizes the luminosity,

$$L_{\text{hot}} \approx 0.15 y^{3/5} \alpha^{7/5} L_{\text{Edd}}, \quad (22)$$

which is obtained by radial integration of eq. (17) ($0.5 L_{\text{hot}}$ comes then from $r \lesssim 40$ and the upper limit of integration is not essential as long as it is $\gg 40$). That case also yields the plasma parameters only weakly changing with r , in particular $\tau \sim 1$ over the hot flow (eq. [16]). Interestingly, the limit of eq. (22) appears to agree with the maximum luminosities observed from black-hole X-ray novae and persistent X-ray binaries in the hard state (Tanaka & Lewin 1995; Tanaka & Shibazaki 1996; Philips et al. 1996; Harmon et al. 1994) and AGNs (Peterson 1997) provided α is large (cf. Narayan 1996; Z98).

The optical depth of the hot flow close to $\dot{M}_{\max}$ is $\tau \sim 1$ (eq. [16]), and $T_e \sim 10^9$ K (eq. [6]), which agree well with the observed X-ray spectra (see Section 1). The values of τ and T_e are basically independent of M (except for a slight dependence via the Coulomb logarithm), which can explain the close similarity of the broad-band spectra of black-hole binaries in the hard state and Seyferts (e.g., Z97, Z98). Also, the solutions very weakly depend on the advection parameter, ξ [e.g., $\tau(\dot{m}_{\max}) \propto \xi^{-1/5}$], which accurate value is obtainable only from global studies. In fact, a more important effect is geometry of Comptonisation models, which should be specified in comparing fit results with hot disc models. E.g., fits in Z97 are for a hemisphere geometry (Poutanen & Svensson 1996), in which case the fitted radial τ is higher than the half-thickness τ obtained from fitting a slab model.

For example, Z98 obtains for GX 339-4 $\tau \simeq 2$ and 0.8 in the hemisphere and slab geometry, respectively. The latter geometry is more appropriate for hot discs radiating close to $\dot{M}_{\text{max}}$, with relatively important cooling.

Black hole systems (in the hard state) show weak spectral variability of the spectral shape with changing flux (see Section 1). We can explain this by the weak dependence of τ (determining the spectral shape) on ℓ_{Coul} . This dependence is $\tau \propto \ell_{\text{Coul}}^{1/6}$ on the cooling-dominated branch, and $\tau \propto \ell_{\text{Coul}}^{2/7}$ on the advection-dominated branch (eqs. [10]–[11]). Which branch corresponds to existing accretion discs can thus be determined by studying the X-ray spectral variability.

The cooling-dominated branch is viscously stable, but unstable thermally (Pringle 1976). We note, however, that this thermal instability is different and much weaker from the classical bremsstrahlung-cooling instability of Pringle, Rees & Pacholczyk (1973), in which an increase of temperature results in a decrease of the cooling rate per unit area. In the model of SLE76, an increase of T_i yields an increase of the cooling rate, but the corresponding increase of the local heating is faster. The flow becomes stable when advection is significant (e.g., Wu 1997).

Finally, we compare the hot solution with the optically-thick (cold) disc solution of Shakura & Sunyaev (1973) and Abramowicz et al. (1988). The right-hand-side curves in Fig. 1 show the latter solution, but with 0.5 of the dissipation occurring in a corona (Svensson & Zdziarski 1994). The solution above $\dot{m} \sim 1$ (weakly dependent on α) has $d\dot{m}/d\tau < 0$ and is both viscously and thermally unstable (Pringle 1981). (The flow becomes again stable due to advection of radiation when $L \sim L_E$, Abramowicz et al. 1988, with L_E corresponding to $\dot{m} = 16$). Thus, the maximum stable $\dot{m} \sim 1$ of the cold solution in Fig. 1 corresponds to $\sim 5\%$ of L_E . Notably, this coincides with the luminosity of Cyg X-1 in the soft state (Gierliński et al. 1998), which supports identification of that state with the lower turning point of the cold solution. (The presence of a corona, assumed above, is inferred from observations of power law tails extending beyond the disc blackbody in the soft state of black-hole binaries, Grove et al. 1998; Gierliński et al. 1998.) We see in Fig. 1 that the $\dot{m}_{\text{max}}$ of the hot solution can be, depending on its α , either larger or smaller than the maximum stable $\dot{m}$ of the cold flow. Thus, the soft-to-hard state transition in X-ray binaries (e.g., Tanaka & Lewin 1995) can in principle correspond to either an increase or decrease of $\dot{M}$.

ACKNOWLEDGEMENTS

This research has been supported in part by the Polish KBN grants 2P03D01008 and 2P03D00614. I thank Marek Gierliński for valuable comments on this work and help with testing the solutions, and Bożena Czerny for discussions and supplying her optically-thick disc code.

REFERENCES

Abramowicz M. A., Chen X., Kato S., Lasota J.-P., Regev O., 1995, *ApJ*, 438, L37
 Abramowicz M. A., Czerny B., Lasota J.-P., Szuszkiewicz E., 1988, *ApJ*, 332, 646

Björnsson G., Abramowicz M. A., Chen X., Lasota J.-P., 1996, *ApJ*, 467, 99
 Chen X., Abramowicz M. A., Lasota J. P., Narayan R., Yi I., 1995, *ApJ*, 443, L61
 Chen X., Abramowicz M. A., Lasota J. P., 1997, *ApJ*, 476, 61
 Chen X., Taam R. E., 1993, *ApJ*, 412, 254
 Dermer C. D., Liang E. P., Canfield E., 1991, *ApJ*, 369, 410
 Esin A. A., 1997, *ApJ*, 482, 400
 Fiore F., Perola G. C., Matsuoka M., Yamauchi M., Piro L., 1992, *A&A*, 262, 37
 Gierliński M., Zdziarski A. A., Coppi P. S., Poutanen J., Ebisawa K., Johnson W. N., 1998, *Nucl. Phys. B*, in press
 Gierliński M., Zdziarski A. A., Done C., Johnson W. N., Ebisawa K., Ueda Y., Haardt F., Philips B. F., 1997, *MNRAS*, 288, 958
 Grove J. E., Johnson W. N., Kroeger R. A., McNaron-Brown K., Skibo J. G., 1998, *ApJ*, 499, in press
 Haardt F., Maraschi L., 1993, *ApJ*, 413, 507
 Harmon B. A., et al., 1994, *ApJ*, 425, L17
 Ichimaru S., 1977, *ApJ*, 214, 840
 Johnson W. N., McNaron-Brown K., Kurfess J. D., Zdziarski A. A., Magdziarz P., Gehrels N., 1997, *ApJ*, 482, 173
 Kuncic Z., Celotti A., Rees, M. J., 1997, *MNRAS*, 284, 717
 Kusunose M., Mineshige S., 1996, *ApJ*, 468, 330
 Liang E. P. T., Thompson K. A., 1979, *MNRAS*, 189, 421
 Magdziarz P., Blaes O. M., Zdziarski A. A., Johnson W. N., Smith D. A., 1998, *MNRAS*, in press
 Mahadevan R., 1997, *ApJ*, 477, 585
 Narayan R., 1996, *ApJ*, 462, 136
 Narayan R., Yi I., 1995, *ApJ*, 452, 710 (NY95)
 Paczyński B., Wiita P. J., 1980, *A&A*, 88, 23
 Peterson B. M., 1997, *Active Galactic Nuclei*, Cambridge University Press, Cambridge
 Philips B. F., et al., 1996, *ApJ*, 465, 907
 Poutanen J., Svensson R., 1996, *ApJ*, 470, 249
 Poutanen J., Svensson R., Stern B., 1997, in Winkler C., Courvoisier T., Durouchoux P., eds., *ESA SP-382, The Transparent Universe*, 401
 Pringle J. E., 1976, *MNRAS*, 177, 65
 Pringle J. E., 1981, *ARA&A*, 19, 137
 Pringle J. E., Rees M. J., Pacholczyk A. G., 1973, *A&A*, 29, 179
 Rybicki G. R., Lightman A. P., 1979, *Radiative Processes in Astrophysics*, Wiley-Interscience, New York
 Sakimoto P. J., Coroniti F. V., 1981, *ApJ*, 247, 19
 Shakura N. I., Sunyaev R. A., 1973, *A&A*, 24, 337
 Shapiro S. L., Lightman A. P., Eardley D. M., 1976, *ApJ*, 204, 187 (SLE76)
 Spitzer, L., 1956, *Physics of Fully Ionized Gases*, Interscience, New York
 Stepney S., Guilbert P. W., 1983, 204, 1269
 Svensson R., Zdziarski A. A., 1994, *ApJ*, 436, 599
 Tanaka Y., Lewin W. H. G., 1995, in: Lewin W. H. G., van Paradijs J., van den Heuvel E. P. J., eds., *X-Ray Binaries*, Cambridge University Press, Cambridge, 126
 Tanaka Y., Shibazaki N., 1996, *ARA&A*, 34, 607
 Ueda Y., Ebisawa K., Done C., 1994, *PASJ*, 46, 107
 Wandel A., Liang E. P., 1991, *ApJ*, 380, 84
 Wu X.-B., 1997, *MNRAS*, 292, 113
 Zdziarski A. A., Johnson W. N., Magdziarz P., 1996, *MNRAS*, 283, 193
 Zdziarski A. A., Johnson W. N., Poutanen J., Magdziarz P., Gierliński M., 1997, in Winkler C., Courvoisier T., Durouchoux P., eds., *ESA SP-382, The Transparent Universe*, 373 (Z97)
 Zdziarski A. A., Poutanen J., Mikołajewska J., Gierliński M., Ebisawa K., Johnson W. N., 1998, *MNRAS*, submitted (Z98)